

## ◇ Traces and Choice ◇

Which traces can be produced by  $P \sqcap Q$  and  $P \sqcap Q$ ? We know that  $P \sqcap Q$  can do the first event of either  $P$  or  $Q$ , and then behave like the remainder of  $P$  or  $Q$ . Therefore any trace of either  $P$  or  $Q$  can be produced by  $P \sqcap Q$ , and we have

$$\text{traces}(P \sqcap Q) = \text{traces}(P) \cup \text{traces}(Q).$$

$P \sqcap Q$  always does  $\tau$  first, and then behaves like either  $P$  or  $Q$ . Because  $\tau$  does not appear in traces, we also have

$$\text{traces}(P \sqcap Q) = \text{traces}(P) \cup \text{traces}(Q).$$

We have previously considered *trace equivalence*, written  $P =_t Q$ , as a definition of when two processes should be considered equal or interchangeable. However, we can now see that  $P \sqcap Q =_t P \sqcap Q$ , even though internal and external choice have been designed to behave in different ways.

In general, trace equivalence is not suitable as a definition of process equivalence.

Before we introduced  $\sqcap$  and  $\sqcup$  all processes were deterministic — the internal state was always determined by the observable events. For deterministic processes, *traces* are all we need to know, and trace equivalence is adequate. But the whole point of introducing the  $\tau$  operator was so that a process could make an internal state change without doing anything observable. Similarly, if  $P$  and  $Q$  have a common event  $a$  available at the first step, then observation of the event  $a$  from  $P \sqcap Q$  does not tell us what the internal state has become.

We will now try to say exactly what the difference between  $P \sqcap Q$  and  $P \sqcup Q$  is, and develop a new notion of process equivalence accordingly.

## ◇ Refusals ◇

Suppose we have the following definitions.

$$P = a \rightarrow P$$

$$Q = b \rightarrow Q$$

What happens if we put each of  $P \sqcap Q$  and  $P \sqcap Q$  in an environment consisting of  $P$ ? i.e. if we look at  $(P \sqcap Q)_{\{a,b\}} \parallel_{\{a,b\}} P$  and  $(P \sqcap Q)_{\{a,b\}} \parallel_{\{a,b\}} P$ .

First, we have  $P \sqcap Q \xrightarrow{a} P$

and  $P \xrightarrow{a} P$

so

$$(P \sqcap Q)_{\{a,b\}} \parallel_{\{a,b\}} P \xrightarrow{a} P_{\{a,b\}} \parallel_{\{a,b\}} P.$$

Also,

$$P_{\{a,b\}} \parallel_{\{a,b\}} P \xrightarrow{a} P_{\{a,b\}} \parallel_{\{a,b\}} P$$

so

$$P_{\{a,b\}} \parallel_{\{a,b\}} P = P$$

(they both satisfy the same recursive definition).

So

$$(P \sqcap Q)_{\{a,b\}} \parallel_{\{a,b\}} P = a \rightarrow P$$

i.e.

$$(P \sqcap Q)_{\{a,b\}} \parallel_{\{a,b\}} P = P.$$

On the other hand,

$$(P \sqcap Q) \{a,b\} \parallel \{a,b\} P \xrightarrow{\tau} P \{a,b\} \parallel \{a,b\} P$$

and

$$(P \sqcap Q) \{a,b\} \parallel \{a,b\} P \xrightarrow{\tau} Q \{a,b\} \parallel \{a,b\} P$$

so

$$(P \sqcap Q) \{a,b\} \parallel \{a,b\} P =$$

$$(P \{a,b\} \parallel \{a,b\} P) \sqcap (Q \{a,b\} \parallel \{a,b\} P).$$

(This is a loose statement as we haven't decided what " $=$ " means yet.)

We know that  $P \{a,b\} \parallel \{a,b\} P = P$

and  $Q \{a,b\} \parallel \{a,b\} P = \text{Stop}$

So

$$(P \sqcap Q) \{a,b\} \parallel \{a,b\} P = P \sqcap \text{Stop}.$$

This shows that  $P \sqcap Q$  and  $P \sqcap Q$  behave differently when put in parallel with  $P$ . One is just  $P$ , the other can internally choose to deadlock (become  $\text{Stop}$ ).

We can use this observation to develop a general approach to distinguishing between nondeterministic processes. We will consider putting a process  $P$  in an environment  $Q$ , where the alphabets of  $P$  and  $Q$  are the same, i.e. constructing  $P \alpha_P \parallel \alpha_P Q$ .

Let  $X$  be a set of events which are offered initially by  $Q$ . If it is possible for  $P \alpha_P \parallel_{\alpha_P} Q$  to deadlock at the first step, then we say that  $X$  is a *refusal* of  $P$ . The set of all refusals of  $P$  is obtained by considering all possible sets  $X$  which could be initial event sets of  $Q$ .

*Examples:* 1. The empty set is a refusal of every process, because if  $Q = Stop$  then  $P \alpha_P \parallel_{\alpha_P} Q = Stop$ .

2. Any set of events  $X$  is a refusal of  $Stop$ .

3. If  $a \notin X$  then  $X$  is a refusal of  $a \rightarrow P$ . So if  $\alpha_P = \{a, b, c\}$  then the refusals of  $a \rightarrow P$  are  $\{\}$ ,  $\{b\}$ ,  $\{c\}$  and  $\{b, c\}$ . Processes  $Q$  causing

$$(a \rightarrow P) \{a,b,c\} \parallel_{\{a,b,c\}} Q$$

to deadlock include  $Stop$ ,  $b \rightarrow Stop$ ,  $c \rightarrow a \rightarrow Stop$ ,  $(b \rightarrow Stop) \square (c \rightarrow c \rightarrow Stop)$ , etc.

4. The refusals of  $(a \rightarrow c \rightarrow Stop) \square (b \rightarrow Stop)$  are  $\{\}$  and  $\{c\}$ .

5. The refusals of  $(a \rightarrow c \rightarrow Stop) \sqcap (b \rightarrow Stop)$  are  $\{\}$ ,  $\{a\}$ ,  $\{b\}$ ,  $\{c\}$ ,  $\{a, c\}$  and  $\{b, c\}$ .

We can define

$$\text{refusals}(P) = \{X \mid X \subseteq \alpha P \text{ and } X \text{ is a refusal of } P\}.$$

Note that  $\text{refusals}(P)$  is a set of sets of events. For example,

$$\text{refusals}((a \rightarrow \text{Stop}) \sqcap (b \rightarrow \text{Stop})) = \{\{\}, \{a\}, \{b\}, \{c\}, \{a, c\}, \{b, c\}\}.$$

In the examples we saw that

$$\text{refusals}((a \rightarrow \text{Stop}) \sqcup (b \rightarrow \text{Stop})) \neq \text{refusals}((a \rightarrow \text{Stop}) \sqcap (b \rightarrow \text{Stop})).$$

In general,  $\text{refusals}(P \sqcup Q) \neq \text{refusals}(P \sqcap Q)$ , and this will be the basis for a new definition of process equality which allows us to distinguish between internal and external choice.

We can now define  $\text{refusals}$  for processes defined in terms of the operators we have seen so far.

$$\text{refusals}(\text{Stop}) = \{X \mid X \subseteq \Sigma\}$$

where  $\Sigma$  is the set of all events being considered — the universal set of events.

$$\text{refusals}(a \rightarrow P) = \{X \mid X \subseteq (\alpha P - \{a\})\}$$

Both of these definitions are subsumed by the definition for menu choice: if  $P = x : A \rightarrow P(x)$  then

$$\text{refusals}(P) = \{X \mid X \subseteq (\alpha P - A)\}$$

If  $P$  can refuse  $X$  then so will  $P \sqcap Q$  if  $P$  is selected. Similarly every refusal of  $Q$  is a possible refusal of  $P \sqcap Q$ .

$$\text{refusals}(P \sqcap Q) = \text{refusals}(P) \cup \text{refusals}(Q)$$

$P \sqcap Q$  can only refuse  $X$  if both  $P$  and  $Q$  can refuse  $X$ .

$$\text{refusals}(P \sqcap Q) = \text{refusals}(P) \cap \text{refusals}(Q)$$

$P \sqcap_A Q$  can refuse all events refused by  $P$  and all events refused by  $Q$ .

$$\text{refusals}(P \sqcap_A Q) = \{X \cup Y \mid X \in \text{refusals}(P) \text{ and } Y \in \text{refusals}(Q)\}$$

Refusals allow us to distinguish formally between deterministic and nondeterministic processes. If a process is deterministic then it can never refuse any event which it could possibly do. In other words, if  $P$  is deterministic and  $a$  is a possible initial event for  $P$ , then  $a$  does not appear in any refusal set of  $P$ .

Writing  $\text{initials}(P)$  for the set of possible initial events of  $P$  (so  $\text{initials}(P) = \{x \mid \langle x \rangle \in \text{traces}(P)\}$ ), we can say that if  $P$  is deterministic then

$$\text{refusals}(P) = \{X \mid X \subseteq \alpha P \text{ and } X \cap \text{initials}(P) = \{\}\}.$$

Determinism means that any event which is possible cannot be taken away by an internal state transition.

Examples. If

$$P = a \rightarrow c \rightarrow Stop \mid b \rightarrow Stop$$

then  $initials(P) = \{a, b\}$  and  $refusals(P) = \{\{\}, \{c\}\}$ .

If

$$P = (a \rightarrow c \rightarrow Stop) \sqcap (b \rightarrow Stop)$$

then  $initials(P) = \{a, b\}$  and (as before)

$$refusals(P) = \{\{\}, \{a\}, \{b\}, \{c\}, \{a, c\}, \{b, c\}\}.$$

Although  $a$  is a possible initial event for  $P$ ,  $P$  could also internally choose to be  $b \rightarrow Stop$  which refuses  $a$ .

To define nondeterminism properly, we need to consider events refused not just at the first step, but after any sequence of events. For example,

$$(a \rightarrow b \rightarrow Stop) \sqcap (a \rightarrow c \rightarrow Stop)$$

is nondeterministic, but this does not become apparent until after the first event.

So:  $P$  is deterministic if and only if

$$\forall s \in traces(P) .$$

$$(refusals(P / s) =$$

$$\{X \subseteq \alpha P \mid X \cap initials(P / s) = \{\}\}).$$

$P / s$  is the process whose behaviour is whatever  $P$  could do after the trace  $s$ .