



STDs: What's New?

The lecturer will address current sexually transmitted disease (STD) management guidelines from the Centers for Disease Control and Prevention (CDC). The lecturer will present a rational approach to cost-effective treatment of patients with STDs and offer a look into future therapeutic options.

- Define a rational, cost-effective approach to STDs based on current CDC recommendations.
- Identify new antibiotics that are of proven efficacy in treating STDs and the additional options for therapy these antibiotics provide.
- Identify antibiotic-resistance patterns of the major STDs and the effects these have on treatment recommendations.

MO-32
Monday, October 11, 1999
1:30 PM - 2:25 PM
Room # N219
Las Vegas Convention Center

*Gregory J Moran, MD, FACEP
Research Grants: Ortho McNeil, Pfizer, Bayer
Speaker: Ortho McNeil, Pfizer, Bayer, Roche

FACULTY

Gregory J Moran, MD, FACEP

Associate Professor, Medicine,
Department of Emergency
Medicine, Division of Infectious
Diseases, Olive View/UCLA Medical
Center, Sylmar, California

Sexually Transmitted Diseases

Gregory J. Moran M.D.

Associate Professor of Medicine

Dept. of Emergency Medicine and Division of Infectious Diseases
Olive View-UCLA Medical Center, Sylmar, CA

Gonorrhea – Treatment Options

Ceftriaxone 125 mg IM

Cefixime 400 mg po

Ciprofloxacin 500 mg po

Ofloxacin 400 mg po

Levofloxacin 250 mg po

Azithromycin 2 gm po

GC - Treatment Options

Cephalosporins:

Treat incubating syphilis

Ceftriaxone 125 mg IM

Best studied at all sites

Requires IM injection

Cefixime 400 mg po

Single dose oral

Equal to Ceftriaxone for cervicitis and urethritis

GC - Treatment Options

Quinolones:

Don't treat incubating syphilis

Contraindicated if pregnant or < 17

Increasing quinolone resistance

Ciprofloxacin 500 mg po - best studied at all sites

Ofloxacin 400 mg po

Effective against Chlamydia if given 7 - 10 days

Levofloxacin 250 mg po

Not FDA approved, but should be equivalent to ofloxacin

GC - Treatment Options

Azithromycin 2 gm po

Single dose for both GC and Chlamydia

GI side effects more common with 2 gm dose

Nausea 18%

Diarrhea 14%

Chlamydia – Treatment Options

Doxycycline 100 mg po bid X 7d

Azithromycin 1 gm po once

Ofloxacin 300 mg bid X 7d

Levofloxacin 250 mg qd X 5d

Erythromycin 500 mg qid X 7d

Antibiotic Compliance - PID

256 women prescribed Doxy 100 bid X 10 d

27% never filled Rx

Reasons: cost 50%

Inconvenience 28%

Sx resolved 20%

42% took incomplete course (mean 3.5 d)


Reasons: Side Effects 36%

Persist. Pain 29%

Sx resolved 22%

Brookoff D: Ann Emerg Med 1992;21: 204

Azithromycin vs. doxy. - Cost



Chlamydia - Diagnosis

Cell culture - Sensitivity 60-90%

Nonculture tests

Direct Fluorescent antibody

Enzyme immunoassay

Nucleic acid hybridization

Polymerase chain reaction

Ligase chain reaction

Chlamydia - Testing

Test	Sensitivity	Specificity
DFA	50-81	88-99
EIA	73-91	95-100
DNA Probe	71-97	97-99

Chlamydia - LCR

Ligase Chain Reaction assay of clean-catch urine vs.
endocervical swab culture

1937 women

	Sensitivity	Specificity
Culture	65	100
Urine LCR	94	99.9

Lee HH. et al. *Lancet* 1995;345:213.

STD vs. UTI

Men - Epididymitis / Prostatitis

Women - Acute Urethral Syndrome

Pyuria without bacteriuria

Cystitis - 10^2 - 10^4 cfu/ml E. coli, etc.

Urethritis - GC, Chlamydia, Herpes

Vaginitis - Trichomonas, Candida

UTI vs. STD

	Doxy	TMP/SMX	Quinolones
Coliforms	+	++	+++
Chlamydia	+++	+	+++
GC	+	+	+++

Pelvic Inflammatory Disease

1 million cases/yr in U.S.

- Ectopic Pregnancy Risk Increased
- Infertility Risk Increased
- Many Subclinical Cases

PID - Diagnosis

Minimal Criteria:

Lower Abdominal Tenderness
Adnexal Tenderness
Cervical Motion Tenderness

Additional Criteria:

Temperature > 38.3
Cervical or Vaginal Discharge
Elevated ESR or C-reactive protein
+ GC or Chlamydia

PID -Admission

Toxic appearing or unable to take po meds
Diagnosis uncertain (cannot r/o appendicitis,
ectopic pregnancy, etc.)
Pelvic abscess suspected
Patient is pregnant
HIV infected
Compliance questionable (e.g. adolescents)
or no follow-up in 72 hrs
Has failed outpatient therapy

PID - Inpatient Rx

Cefoxitin 2 gm q6 or Cefotetan 2 gm q12
plus Doxycycline 100 mg q 12

or

Clindamycin 900 mg q 8
plus Gentamicin

or

Azithromycin 500 mg IV qd X 1-2 days,
then 250mg qd to complete 7 days
plus Metronidazole

PID - Outpatient Rx

Ofloxacin 400 + Metronidazole 500 bid x 14d

or

Ceftriaxone 250mg IM + Doxy 100 bid x 14 d

STD - Partner Treatment

168 women with Chlamydia infection

125 given referral cards for partner

43 given doxycycline for partner

Rate of Chlamydia recurrence over 18 mos

Card: 22. 1% per year

Doxy: 11. 5% per year (OR 0.37)

Kissinger P. *Sex Trans Inf* 1998;74:331.

Trichomonas - Treatment

Metronidazole

2gm po single dose

or 500mg bid X 7d

Safe in pregnancy

Topical preparations not effective

Chancroid

Haemophilus ducreyi

Therapy:

Azithromycin 1 gm orally
Ceftriaxone 250mg IM
Ciprofloxacin 500 bid X 3d
Erythromycin 500mg qid X 7d

Herpes simplex - Treatment

First episode

Acyclovir 400 tid or 200 5X/d for 7-10 days
Famcyclovir 250 mg tid for 5-10 days
Valacyclovir 1 gm bid for 7-10 days

Recurrent episodes

Acyclovir 400 mg tid for 5 days
Famcyclovir 125 mg bid for 5 days
Valacyclovir 500 mg bid for 5 days

ED Screening for Syphilis

Should test ALL pts. treated for STD

Often overlooked:

Frequency of syphilis testing in GC pts:

Males 58%

Females 26%

Frequency of + VDRL/FTA

in pts. with STD complaints: 6%

Kirsch TD: Ann Emerg Med (abstract)1992; 21: 204
Ernst A: Ann Emerg Med 1991; 20:

Syphilis - Serologic Tests

Nontreponemal - VDRL or RPR

Quantitative - Rise with new infection and fall after treatment

Cheaper – used for screening

May be false positive

Treponemal - FTA ABS or MHATP

Qualitative + or -

Once positive, always positive

More specific

Syphilis - Therapy

Primary, Secondary, Early Latent < 1 year

Benzathine PCN 2.4 million units IM X 1

or Doxycycline 100mg bid X 2 wks

Late Latent > 1 year, Gummas, Cardiovascular

Benzathine PCN G 2.4 million units weekly X 3

or Doxycycline 100mg bid X 4 weeks

Syphilis - Lumbar Puncture

All patients with neurologic signs

Syphilis > 1 year's duration with:

Treatment failure or non-PCN Rx planned

VDRL \geq 1:32 or HIV +

Other evidence of active syphilis

Neurosyphilis Therapy:

PCN G 2-4 million units q4h IV X 10-14d

Sexual Assault

Evaluation:

GC, Chlamydia, VDRL, pregnancy
HIV, Hep B

Prophylaxis:

Cefixime 400 mg po (or alternative GC Rx)
Azithromycin 1 gm po (or doxy 100 bid X 7d)
Metronidazole 2 gm po
Hepatitis B - begin vaccine series
? HIV

CDC. *MMWR* 1998;47(RR-1)

HIV Prophylaxis for Sexual Exposure

Risk for receptive anal intercourse > needlestick

Risk for receptive vaginal intercourse < needlestick
- approximately 1 /1 000

Regimen: AZT 300mg bid or 200 mg tid
plus 3TC 150 mg bid
for 4 weeks

Consider adding nelfinavir 750 tid or indinavir 800 tid
for highest risk exposures

Bamberger JD et al., *Am J Med* 1999;106:323

Selected Reading - STDs

CDC: 1998 Sexually Transmitted Disease Treatment Guidelines. *MMWR* 1998;47(RR-1):1-116.

Moran GJ. Diagnosing Sexually Transmitted Diseases-Ulcerating Diseases. *Emergency Medicine* 1997; 29:58-72.

Moran GJ. Diagnosing and Treating STDs - Lesionless Disorders. *Emergency Medicine* 1997; 29: 20-3 1.

Eng TR, Butler WT, eds. The Hidden Epidemic:Confronting STDs. National Academy Press, Wash., DC, 1997.

Moran JS, Levine WC. Drugs of choice for the treatment of uncomplicated gonococcal infections. *Clin Infect Dis* 1995; 20(Supp 1): S47-65.

Weber JT, Johnson RE. New treatments for *Chlamydia trachomatis* genital infection. *Clin Infect Dis* 1995; 20(Supp 1): S66-71.

CDC. Fluoroquinolone-resistant *Neisseria gonorrhoeae*--San Diego, CA, 1997. *MMWR* 1998,47(20):405-8.

- Haddix AC, Hillis SD, Kassler WJ. The cost effectiveness of azithromycin for *Chlamydia trachomatis* infections in women. *Sex Trans Dis* 1995; 22: 274-80.
- Woods GL. Update on laboratory diagnosis of STDs. *Clin Lab Med* 1995; 15: 665-684.
- McCormack WM: Current Concepts: Pelvic Inflammatory Disease. *New Engl J Med* 1994; 330: 115-19
- Hemsell DL, Little BB, Faro S, et al. Comparison of three regimens recommended by the CDC for the treatment of women hospitalized with acute PID. *Clin Infect Dis* 1994; 19: 720-7.
- Centers for Disease Control: Pelvic Inflammatory Disease: Guidelines for Prevention and Management. *MMWR* 1991;40(supp RR-5): 1.
- Whitley RJ, Kimberlin DW, Roizman B. Herpes Simplex Viruses. *Clin Infect Dis* 1998; 26: 541-553.
- Wald A. New therapies and prevention strategies for genital herpes. *Clin Infect Dis* 1999;28(Supp1):S4-S13.
- Augenbraun MH; McCormack WM. Sexually transmitted diseases in HIV-infected persons. *Infect Dis Clin North Am*, 1994 Jun, 8(2):439-48.
- Martin DH, Mroczkowski TF, Dalu ZA et al. A controlled trial of a single dose of azithromycin for the treatment of chlamydial urethritis and cervicitis. *New Engl J Med* 1992; 327: 921-5.
- Augenbraun MH, Rolfs R. Treatment of syphilis, 199X. *Clin Infect Dis* 1999;28(Supp1):S21-S28.
- Beutner KR, Wiley DJ, Douglas JM, et al. Genital warts. *Clin Infect Dis* 1999;28(Supp1):S37-S56.
- Joesoef MR, Schmid GP, Hillier SL. Bacterial vaginosis. *Clin Infect Dis* 1999;28(Supp1):S57-S65.
- Jenny C; Hooton TM; Bowers A, Copass MK; Krieger JN; Hillier SL; Kiviat N; Corey L; Stamm WE; Holmes KK. Sexually transmitted diseases in victims of rape. *New Engl J Med*, 1990 Mar 15, 322(11):713-6.
- Beck-Sague CM, Solomon F. Sexually transmitted diseases in abused children and adolescent and adult victims of rape: Review of selected literature. *Clin Infect Dis* 1999;28(Supp1):S74-S83.
- Royce RA, Sefia A, Cates W, Cohen MS. Sexual transmission of HIV *NEJM* 1997; 336: 1072-S.
- CDC. Management of possible sexual, injecting-drug-use, or other nonoccupational exposure to HIV, including considerations related to antiretroviral therapy. *MMWR* 1998;47(RR-17): 1-14.
- Katz MH, Gerberding JL. The care of persons with recent sexual exposure to HIV, *Ann Intern Med* 1998;128:306-312.
- Lurie P, Miller S, Hecht F, Chesney M, Lo B. Postexposure prophylaxis after nonoccupational HIV exposure. *JAMA* 1998;280:1769-1773.
- Bamberger JD, Waldo CR, Gerberding JL, Katz MH. Postexposure prophylaxis for human immunodeficiency virus infection following sexual assault. *Am J Med* 1999;106:323-6.

ATT09889.txt

Gregory J. Moran MD
Assistant Professor of Medicine tel 818 364 3110
UCLA School of Medicine fax 818 364 3268
Dept. of Emergency Medicine
and Division of Infectious Diseases
Olive View - UCLA Medical Center
14445 Olive View Drive
Sylmar, CA 91342 email gmoran@ucla.edu
