

# hairy woodpecker

Picoides villosus

Kingdom: Animalia
Division/Phylum: Chordata
Class: Aves

Order: Piciformes Family: Picidae

### **FEATURES**

The hairy woodpecker averages nine and one-half inches in length. Its back, outer tail feathers, belly, stripes on its head and spots on its wings are all white. The head, wings and tail are black. The male has a small, red patch on the back of the head. Unlike the similar downy woodpecker, the hairy woodpecker has a large bill.

#### **BEHAVIORS**

The hairy woodpecker is a common, permanent resident statewide. Nesting takes place from March through June. The nest cavity is dug five to 30 feet above the ground in a dead or live tree. Both sexes excavate the hole over a one- to three-week period. The female deposits three to six white eggs on a nest of wood chips. The male and female take turns incubating the eggs during the day, with only the male incubating at night for the duration of the 11- to 12-day incubation period. One brood is raised per year. The hairy woodpecker lives in woodlands, wooded areas in towns, swamps, orchards and city parks. Its call is a loud "peek." This woodpecker eats insects, seeds and berries.

### **HABITATS**

- bottomland forests
- coniferous forest
- southern Illinois lowlands
- upland deciduous forest

## **ILLINOIS STATUS**

- common endangered threatened

# **ILLINOIS RANGE**

Permanent resident: statewide

Summer resident:

Migrant:

Winter resident:

#### **BIBLIOGRAPHY**

Illinois Department of Natural Resources. 1999. Biodiversity of Illinois, Volume 2: Woodland Habitats CD-ROM.