

# black vulture

Coragyps atratus

- Kingdom: Division/Phylum: Chordata Class: Order: Family:
  - Animalia Aves Falconiformes Cathartidae

### **FEATURES**

The black vulture averages 23 to 27 inches in length. It has a short tail. Except for white patches in the wing tips which can be seen as it flies, this bird is entirely black, including its bare head and legs.

### **BEHAVIORS**

The black vulture is an uncommon, permanent resident in extreme southern Illinois. It is occasionally seen as a vagrant elsewhere in the state. Nesting occurs in April and May. No nest is built. The female lays two eggs in a shallow cave or tree cavity. The gray-green to dull white eggs have brown marks. Both the male and female take turns incubating during the 28- to 39-day incubation period. This bird lives in swampy woodlands and bluffs. It feeds by scavenging for dead organisms.

### HABITATS

bottomland forests coniferous forest southern Illinois lowlands upland deciduous forest

## **ILLINOIS STATUS**

common native

endangered exotic

threatened

### **ILLINOIS RANGE**

Permanent resident: southern tip Summer resident: Migrant: Winter resident:

#### **BIBLIOGRAPHY**

Illinois Department of Natural Resources. 1999. Biodiversity of Illinois, Volume 2: Woodland Habitats CD-ROM.